Second law of thermodynamics with quantum memory

Li-Hang Ren[1,2] and Heng Fan[1,2,3,*]

[1]*Beijing National Laboratory for Condensed Matter Physics, Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*
[2]*School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100190, China*
[3]*Collaborative Innovation Center of Quantum Matter, Beijing 100190, China*

We design a heat engine with multi-heat-reservoir, ancillary system and quantum memory. We then derive an inequality related with the second law of thermodynamics, and give a new limitation about the work gain from the engine by analyzing the entropy change and quantum mutual information change during the process. In addition and remarkably, by considering two measurements and with the help of the entropic uncertainty relation with quantum memory, we find that the total work gains from the heat engine should be larger than a quantity related with quantum entanglement between the ancillary state and the quantum memory. This result provides a lower bound for the work extracted, in contrast with the upper bound in the conventional second law of thermodynamics, and thus shed light from an opposite direction for the foundations of the thermodynamics.

PACS numbers: 03.67.-a, 05.70.-a, 89.70.Cf, 03.65.Ta

arXiv:1701.07628v1 [quant-ph] 26 Jan 2017

Introduction.— Maxwell's demon plays an important role in the history of thermodynamics and information theory [1, 2]. It is first proposed by Maxwell that a powerful demon might conduct microscopic operation to break the second law of thermodynamics. However, according to Landauer's principle, manipulation of information will be accompanied by energy consumption [3], which saves the second law, see also [4]. However, with this demon, one can relax the restrictions imposed by the second law on the energy exchanged between a system and surroundings, and some new thermodynamic inequalities are studied, see [5] and a review [6]. In conventional thermodynamics, the second law gives $W_{ext} \leq -\triangle F$, where W_{ext} is the extractable work from system and $F = U - TS$ is the free energy during the isothermal process. Due to the Maxwell's demon, this thermodynamic expression can be extended to a favorable form with discrete quantum feedback control[7, 8]:

$$W_{ext} \leq -\triangle F + k_B T I, \tag{1}$$

in which k_B is the Boltzmann constant and I is the quantum-classical mutual information describing the mutual information of a fixed quantum system and the outcome classical information obtained by a quantum measurement. This quantum-classical mutual information is an extension of the standard quantum mutual information defined originally between two subsystems. One may then observe that the maximum work that can be extracted may exceed that in conventional thermodynamics, however, the marginal part is restricted by term of the quantum-classical mutual information. This inequality lays an extension of the second law of thermodynamics.

The above statement shows information can be exploited to extract physical work, which may be called an information heat engine [9]. Szilard first proposed a physical model, i.e. the so-called Szilard engine (SZE), to realize Maxwell's demon [10]. This SZE involves a single-molecule gas in a box, immersed in a thermal reservoir at temperature T, and an external demon. The demon inserts a partition into the middle of the box, then measures on which side the molecule is trapped and performs expansion to extract work $W_{ext} = k_B T \ln 2$. In SZE, the information that the molecule is in the left or right is exploited to extract physical work. Both theoretical and experimental studies on the information heat engine, additionally the extension for quantum case, have been performed [7–20]. Quantum resources for quantum information processing such as quantum entanglement, quantum discord or quantum coherence may probably be converted to extractable work. It is proved that the work gain may result from the entanglement between subsystems [11] because of deep connections between thermodynamics and the theory of entanglement [12, 13]. Also one can devise a heat engine which can be driven by purely quantum information such as the quantum discord [9]. Recently, experimental investigation is performed to show that quantum discord is necessary in energy transport in a nanoscale aluminum-sapphire interface [14].

We know that for a quantum system, a quantum memory can be available and quantum entanglement or some quantumness of correlations may play a critical role in the emergent quantum phenomena [21], such as that in the entropic uncertainty relation with a quantum memory [22, 23] generalizing Heisenberg uncertainty principle [24]. As we may notice that entanglement and quantum discord appear to be resource for work extraction in thermodynamics. It is desirable to construct a heat engine where quantum correlations or entanglement appear involving in the process. In this Letter, we design a heat engine which includes the system S contacted by independent heat baths with possible different temperatures, the ancillary system A and a quantum memory B, see Fig. 1. This set up of heat engine is similar with that in Ref.[7], but with an ancillary system A and a quantum memory B. With the help of the quantum memory, we then can characterize the role of quantum correlation in the thermodynamic circle. Thus, new second law of thermodynamic inequality can be obtained.

Set up of the heat engine and its process.— The heat engine involves four parts: system S, a set of thermal reservoirs $R = \{R_1, \cdots R_n\}$ and composite quantum system M consisting

*Electronic address: hfan@iphy.ac.cn



FIG. 1: Set up of the heat engine. The system S contacts with independent heat baths $R_1, R_2, ..., R_n$ at temperatures $T_1, T_2, ..., T_n$ respectively constituting the multi-heat-reservoir R, the system S has ancillary system A, where we can make measurement. Particularly, the set up includes a quantum memory B which is possibly entangled with ancillary state A. At stage (ii), a unitary operation $U_{SR}^{(1)}$ is performed globally on the system and the multi-heat-reservoir SR.



FIG. 2: Processing of the heat engine. In the thermodynamic process, at stages (iii) and (iv), a unitary transformation $U^{(2)}$ is performed on system and ancillary state SA, followed by a projective measurement on ancillary system A. Depending on measurement results k, a feedback control unitary operation U_{SR}^{k} is then implemented on the system and the multi-heat-reservoir SR.

of ancillary state A and quantum memory B, see Fig. 1. The total Hamiltonian is written as

$$H(t) = H_{SR}(t) + H_{SM_{AB}}^{int}(t) + H_{M_{AB}}(t), \qquad (2)$$

where $H_{SR}(t) = H_S(t) + \sum_{m=1}^{n}[H_{SR_m}^{int}(t) + H_{R_m}]$, describing the Hamiltonian of the system, reservoirs and their interaction. During the operating process, system S can contact $R_1, R_2, \ldots, R_n$ which are at respective temperatures $T_1, T_2, \ldots, T_n$. By contacting S with R_1 at the start and the end of the process, system S is in thermodynamic equilibrium in the initial and final state. However, the system may not be in thermodynamic equilibrium between the initial and final state. For convenience, we note the temperature of S in the initial and final state as $T = T_1$. At the beginning and last, we assume $H_{SR_m}^{int}(t_i) = H_{SR_m}^{int}(t_f) = 0$, $H_S(t_i) = H_S^{(i)}$ and $H_S(t_f) = H_S^{(f)}$. The general process of the engine is divided into four stages, which is similar as that in Ref.[7] but with different operations since we have additional quantum systems A and B.

Stage (i). At time t_i, the system and reservoirs are in thermodynamic equilibrium respectively, that is, they are in canonical distribution. The initial state reads

$$\begin{aligned}\rho^{(i)} &= \frac{\exp(-\beta H_S^{(i)})}{Z_S^{(i)}} \otimes \frac{\exp(-\beta_1 H_{R_1})}{Z_{R_1}} \otimes \cdots \\ &\otimes \frac{\exp(-\beta_n H_{R_n})}{Z_{R_n}} \otimes \rho_{AB}^{(i)} \\ &\equiv \rho_{SR}^{(i)} \otimes \rho_{AB}^{(i)} \end{aligned} \qquad (3)$$

where $\beta_n = 1/(k_B T_n)$ is related with temperature T_n, and the partition functions are $Z_S^{(i)} = \mathrm{Tr}[\exp(-\beta H_S^{(i)})]$ and $Z_{R_n} = \mathrm{Tr}[\exp(-\beta_n H_{R_n})]$.

Stage (ii). System S begins to interact with the surrounding to extract work. In a general way any thermodynamic process between the system and reservoirs can be expressed by an unitary evolution. With unitary operation $U^{(1)} = I_{AB} \otimes U_{SR}^{(1)}$ as shown in Fig. 1, the initial state becomes

$$\rho^{(1)} = U^{(1)} \rho^{(i)} U^{(1)\dagger}, \qquad (4)$$

in which I_{AB} is the identity operator for M consisting of ancillary state A and memory B. During this process, the composite system M is left unchanged.

In stage (iii), the system M starts to work, which plays the role of Maxwell's demon. In order to make use of quantum information to extract work, we let the system S interacts with the ancillary system A such that there is quantum information exchange, then we make measurement on A by positive operator valued measures (POVMs). Specifically speaking, this measurement process is performed by A with rank-1 projector $\{\Pi_A^k\}$. So the measurement process is implemented by performing a unitary transformation $U^{(2)}$ on SA followed by a projection measurement $\{\Pi_A^k : |k\rangle\langle k|\}$ on A. The density matrix is given by

$$\begin{aligned}\rho^{(2)} &= \sum_k \Pi_A^k U^{(2)} \rho^{(1)} U^{(2)\dagger} \Pi_A^k \\ &= \sum_k p_k |k\rangle_A\langle k| \otimes \rho_{BSR}^{(2)k} \end{aligned} \qquad (5)$$

The measurement outcome $p_k = tr[\Pi_A^k U^{(2)} \rho^{(1)} U^{(2)\dagger} \Pi_A^k]$ is registered by the memory and the post measurement state is $\rho_{BSR}^{(2)k} = tr_A[\Pi_A^k U^{(2)} \rho^{(1)} U^{(2)\dagger} \Pi_A^k / p_k]$.

The stage (iv) is the feedback control. It is performed discretely depending on outcome p_k by applying corresponding operations on the system S and the multi-heat-reservoir R. Feedback control is also a quantum process, the unitary operator is written as $U^{(3)} = I_B \otimes \sum_k |k\rangle_A\langle k| \otimes U_{SR}^k$. The whole process of stage (iii) and the feedback control of stage (iv) are schematically presented in Fig. 2. Now, the final state becomes

$$\rho^{(f)} = U^{(3)} \rho^{(2)} U^{(3)\dagger}. \qquad (6)$$

In the last equilibration process, the system and heat reservoirs evolve to reach thermodynamic equilibrium, which can be described by a unitary transformation. As unitary transformation has no influence on the following calculation. Thus

we ignore this process, and regard $\rho^{(f)}$ as the final state. Although the system and reservoirs are equilibrium states at last, they may not equal the canonical distribution. In order to evaluate the energy of the system, we introduce the canonical distributed state as a reference which takes the standard form,

$$\rho_{SR}^{(ref)} = \frac{\exp(-\beta H_S^{(f)})}{Z_S^{(f)}} \otimes \frac{\exp(-\beta_1 H_{R_1})}{Z_{R_1}} \otimes \cdots \otimes \frac{\exp(-\beta_n H_{R_n})}{Z_{R_n}}, \quad (7)$$

where $Z_S^{(f)} = \mathrm{Tr}[\exp(-\beta H_S^{(f)})]$. The reference state will be used to correspond a equilibrium state which may not be in the canonical distribution form.

Maximum work extracted from the heat engine.— We will proceed to calculate the net work gain from the heat engine by analyzing the entropy change during the process. The von Neumann entropy of the state ρ is defined as $S(\rho) = -\mathrm{Tr}(\rho \ln \rho)$. The following discussion will involve the technique of quantum information [21].

The difference between states $\rho^{(i)}$ and $\rho^{(1)}$ is a unitary transformation, so the entropy remains invariant. However, in the third step, measurement is performed. Thus, we know that projective measurements increase entropy, one obtains

$$S[\rho^{(i)}] \leq S[\rho^{(2)}]. \quad (8)$$

According to equalities (3) and (5), we know that states $\rho^{(i)}$ is product state, $S[\rho^{(i)}] = S[\rho_{SR}^{(i)}] + S[\rho_{AB}^{(i)}]$, also state $\rho^{(2)}$ can be written as the following form,

$$S[\rho^{(2)}] = H(p_k) + \sum_k p_k S[\rho_{BSR}^{(2)k}]. \quad (9)$$

With the help of the subadditivity of von Neumann entropy, $S[\rho_{BSR}^{(2)k}] \leq S[\rho_{SR}^{(2)k}] + S[\rho_B^{(2)k}]$, and the above facts, the initial inequality (8) now takes the form,

$$S[\rho_{SR}^{(i)}] - \sum_k p_k S[\rho_{SR}^{(2)k}] \leq H(p_k) + \sum_k p_k S[\rho_B^{(2)k}] - S[\rho_{AB}^{(i)}] \quad (10)$$

By some calculations, see supplementary material for detail [25], we can find the entropy change between initial state and the final state for system S and the multi-heat-reservoir R,

$$S[\rho_{SR}^{(i)}] - S[\rho_{SR}^{(f)}] \leq \triangle S_A + \triangle S_B - \triangle I, \quad (11)$$

where $\triangle S_A = S[\rho_A^{(f)}] - S[\rho_A^{(i)}]$ denotes the entropy change for ancillary system A, and $\triangle S_B$ denotes the entropy change of the quantum memory B, I denotes quantum mutual information defined as $I(X:Y) = S(\rho_X) + S(\rho_Y) - S(\rho_{XY})$, here $\triangle I = I(A^{(2)} : B^{(2)}) - I(A^{(i)} : B^{(i)})$ represents the change of quantum mutual information in the process of heat engine for composite system M including both A and B.

According to Klein's inequality $S(\rho\|\sigma) = tr(\rho \ln \rho) - tr(\rho \ln \sigma) \geq 0$, we have $tr[\rho_{SR}^{(f)} \ln \rho_{SR}^{(ref)}] \leq -S[\rho_{SR}^{(f)}]$. Therefore,

$$S[\rho_{SR}^{(i)}] + tr[\rho_{SR}^{(f)} \ln \rho_{SR}^{(f)can}] \leq \triangle S - \triangle I \quad (12)$$

with $\triangle S \equiv \triangle S_A + \triangle S_B$. From Eq.(3) and Eq.(7), we know that $\rho_{SR}^{(i)}$ and $\rho_{SR}^{(ref)}$ are canonical distribution. Then substituting their specific expression into inequality (11), we obtain

$$E_S^{(i)} - E_S^{(f)} + \sum_{m=1}^{n} \frac{T}{T_m}(E_{R_m}^{(i)} - E_{R_m}^{(f)}) \leq F_S^{(i)} - F_S^{(f)} + k_B T[\triangle S - \triangle I], \quad (13)$$

where $E_S^{(i)} = \mathrm{Tr}(\rho^{(i)} H_S^{(i)})$, $E_{R_m}^{(i)} = \mathrm{Tr}(\rho^{(i)} H_{R_m})$, $F_S^{(i)} = -k_B T \ln Z_S^{(i)}$, $E_S^{(f)} = \mathrm{Tr}(\rho^{(f)} H_S^{(f)})$, $E_{R_m}^{(f)} = \mathrm{Tr}(\rho^{(f)} H_{R_m})$ by comparing the final state to the reference state, also $F_S^{(f)} = -k_B T \ln Z_S^{(f)}$. Among them $E_S^{(f)} - E_S^{(i)} \equiv \triangle U_S$ is the change of the internal energy, $E_{R_m}^{(i)} - E_{R_m}^{(f)} \equiv Q_m$ is the heat exchange between system and reservoir R_m and $F_S^{(f)} - F_S^{(i)} \equiv \triangle F_S$ is the difference in the Helmholtz free energy of system. Then the above inequality becomes

$$-\triangle U_S + \sum_{m=1}^{n} \frac{T}{T_m} Q_m \leq -\triangle F_S + k_B T[\triangle S - \triangle I]. \quad (14)$$

This inequality is one of the main results in this Letter. Before interpreting this extension of the second law of thermodynamic inequality in more detail, two special cases will first be illustrated.

If $n = 1$, there is only one reservoir with temperature T. As the work extractable from the engine is defined as $W_{ext} = -\triangle U_S + Q = -(E_S^{(f)} - E_S^{(i)}) + E_R^{(i)} - E_R^{(f)}$, the final result (14) reduces to a simple case

$$W_{ext} \leq -\triangle F_S + k_B T \triangle S - k_B T \triangle I. \quad (15)$$

This inequality is in agreement with the result of Ref. [9] where the quantum-classical mutual information is introduced. Here we use the standard quantum mutual information. The inequality means that the extracted work can exceed the difference of the free energy which generalizes the conventional second law of thermodynamics, however, the marginal part is constrained by the difference of the changes for entropy and the quantum mutual information $k_B T(\triangle S - \triangle I)$.

When $n = 2$, the heat engine becomes an analogue Carnot cycle. We take two heat baths for consideration: $T_H > T_L$. After a cycle, we assume that $\triangle U_S = \triangle F_S = 0$. Because $W_{ext} = Q_H + Q_L$, we find,

$$W_{ext} \leq \left(1 - \frac{T_L}{T_H}\right) Q_H + k_B T_L(\triangle S - \triangle I) \quad (16)$$

The efficiency of the heat engine is

$$\eta = \frac{W_{ext}}{Q_H} = 1 - \frac{T_L}{T_H} + \frac{k_B T_L(\triangle S - \triangle I)}{Q_H}. \quad (17)$$

In contrast with the efficiency of the conventional Carnot cycle $\eta_{carnot} = 1 - \frac{T_L}{T_H}$, the heat engine presented here can exceed the conventional Carnot heat engine, but new restrictions is still imposed as presented in the last term.

The general n case heat engine can be considered as simple extension for $n = 1, 2$, and the inequality can be considered as

the second law of thermodynamics with quantum correlation in quantum information science. We comment that the mutual information can be divided into two parts as classical correlation and quantum discord [26], so $\triangle I = \triangle J + \triangle\delta$. From the Eq.(5), we have $\rho_{AB}^{(2)} = \sum_k p_k|k\rangle_A\langle k|\otimes\rho_B^{(2)k}$ which is a post-measurement density matrix, so discord $\delta(B^{(2)}|A^{(2)}) = 0$. Thus for example $n = 1$, Eq.(15) becomes

$$W_{ext} \leq -\triangle F_S + k_B T C, \tag{18}$$

where we use the notation, $C = \triangle S - \triangle J + \delta(B^{(i)}|A^{(i)})$. It shows that the initial quantum discord can be exploited to acquire physical work, in agreement with the results in [14]. Similar form can also be obtained for general n from Eq.(14).

Lower bound for work gained with different measurement bases.— The heat engine in this work includes measurement process at the third stage. It is well-known that Heisenberg has asserted a fundamental limit to the precision of the outcomes for a pair of incompatible obervables [24]. For a quantum system probably entangled with a quantum memory, there exists the entanglement-assisted entropic uncertainty relation for two incompatible measurements [22], and more generally for multiple measurements [23]. Next, we will study the heat engine, however, by considering two measurements at stage (iii). For convenience, we just consider the single-reservoir case.

We emphasize that we next concentrate on the measurement process. The heat engine will work twice with two sets of measurement operators, $\{\Pi_A^k : |k\rangle\langle k|\}$ and $\{\Pi_A^m : |m\rangle\langle m|\}$. We denote the maximal overlap of the two sets of projective operators as, $c = \max_{k,m}|\langle k|m\rangle|^2$. For state ρ_{AB}^i, we have the entropic uncertainty relation,

$$S(k|B) + S(m|B) \geq \log_2\frac{1}{c} + S(A|B), \tag{19}$$

in which $S(A|B) = S(\rho_{AB}^i) - S(\rho_B^i)$ is the conditional entropy for initial state ρ_{AB}^i, we would like to point out that $S(A|B)$ can be negative for entangled initial state. The quantity $S(k|B)$ is the conditional von Neumann entropy of the post-measurement state after performing the measurement $\{\Pi_A^k\}$ on A,

$$S(k|B) = S[\sum_k p_k|k\rangle\langle k|\otimes\rho_B^{(i)k}] - S(\rho_B^i), \tag{20}$$

where $p_k = tr(\Pi_A^k\rho_{AB}^i\Pi_A^k)$ and $\rho_B^{(i)k} = tr_A(\Pi_A^k\rho_{AB}^i\Pi_A^k)/p_k$. Similarly, $S(m|B)$ can also be defined for measurement $\{\Pi_A^m\}$ as follows: $S(m|B) = S[\sum_m q_m|m\rangle\langle m|\otimes\rho_B^{(i)m}] - S(\rho_B^i)$, where $q_m = tr(\Pi_A^m\rho_{AB}^i\Pi_A^m)$ and $\rho_B^{(i)m} = tr_A(\Pi_A^m\rho_{AB}^i\Pi_A^m)/q_m$. We then substitute these equalities into relation (19), the outcomes should satisfy the uncertainty relation.

By considering the working process of the heat engine, we know that $\rho_{AB}^{(2)} = Tr_{SR}\rho^{(2)} = \sum_k p_k|k\rangle_A\langle k|\otimes\rho_B^{(2)k}$. For partial trace, $\rho_B^{(i)k} = tr_A(\Pi_A^k\rho_{AB}^i\Pi_A^k)$ and $\rho_B^{(2)k} = tr_{SR}[tr_A(\Pi_A^k\rho^{(1)\prime}\Pi_A^k)] = tr_A(\Pi_A^k\rho_{AB}^i\Pi_A^k)$. Then we have $\rho_B^{(i)k} = \rho_B^{(2)k}$. Thus the uncertainty relation (19) can be rewritten as,

$$S(\rho_{AB}^{(2)}) + S(\rho_{AB}^{(2)\prime}) \geq \log_2\frac{1}{c} + S(\rho_{AB}^i) + S(\rho_B^i), \tag{21}$$

where $\rho_{AB}^{(2)\prime}$ indicates the corresponding state when we use the measurement bases of $\{\Pi_A^m\}$.

The extractable work has been illustrated in Eq.(15). Note that the relation (11) gives $\triangle S - \triangle I = S[\rho_{AB}^{(2)}] - S[\rho_{AB}^{(i)}]$. Next we denote W_{max}^k as the maximum extractable work of the first engine with projector $\{\Pi_A^k : |k\rangle\langle k|\}$, and the similar as W_{max}^m for another measurement. Then

$$W_{\max}^k = -\triangle F_S + k_B T[S(\rho_{AB}^{(2)}) - S(\rho_{AB}^{(i)})], \tag{22}$$

$$W_{\max}^m = -\triangle F_S' + k_B T[S(\rho_{AB}^{(2)\prime}) - S(\rho_{AB}^{(i)})]. \tag{23}$$

Due to the limitation of the entropic uncertainty relation, $S(\rho_{AB}^{(2)}) + S(\rho_{AB}^{(2)\prime})$ satisfy the inequality (21). We put it into the relation $W_{\max}^k + W_{\max}^m$. Then the final result reads

$$W_{\max}^k + W_{\max}^m \geq -\triangle F_S - \triangle F_S' + k_B T[\log_2\frac{1}{c} - S(A^i|B^i)]. \tag{24}$$

This inequality is also one of our main results. The term $S(A^i|B^i)$ appearing on the right-hand side quantifies the amount of entanglement between A and B. We remark that the quantity $S(A^i|B^i)$ plays a significant role in quantum information science, see for example [27–29] and the references therein.

The inequality (24) plays a complete different role in comparing with the inequalities of second law of thermodynamics (14-16). Remarkably, the entropic uncertainty which is a representation of the Heisenberg uncertainty principle implies that there exists lower bound, instead of upper bound as shown in (14-16), for the heat engine presented in this Letter. That is to say, the total work gain can be larger than a bound for two measurement bases. The negative $S(A^i|B^i)$, meaning the existence of entanglement, will result in higher lower bound and let the work extracted larger for two measurements. This is a new phenomenon which is not observed before. Straightforwardly, our results can be generalized for multiple measurements.

Conclusion.—We design a specific heat engine with both ancillary state and the quantum memory. The new inequality related with the second law of thermodynamics is obtained. For simple cases, our results extend the results in the isothermal process and the Carnot circle. The changes of entropy and the quantum mutual information lay new limit for the marginal part of work which exceeds the conventional second law of thermodynamics. Surprisingly, if two measurements are preformed, we find a new inequality due to the entropic uncertainty relation with the assistance of quantum memory, which provides a lower bound for the work gained from the heat engine. This result describes an opposite fact of the thermodynamics by considering entanglement in quantum information science. We believe that our results offer new critical ingredients for the foundations of quantum thermodynamics.

Acknowledgments: This work was supported by MOST of China (Grants No. 2016YFA0302104 and

2016YFA0300600), national natural science foundation of China NSFC (Grant No. 91536108) and Chinese Academy of Sciences (Grants No. XDB01010000 and XDB21030300). We thank Zheng-An Wang for helping us in drawing the pictures.

[1] J. C. Maxwell, *Theory of Heat*(Appleton, Lonton, 1871).
[2] H. S. Leff and A. F. Rex, *Maxwell's demons 2*(IOP Publishing, Bristol, 2003).
[3] R. Landauer, IBM J. Res. Dev. **5**, 183 (1961).
[4] C. H. Bennett, Int. J. Theor. Phys. **21**, 905 (1982).
[5] S. Lloyd, Phys. Rev. A **56**, 3374 (1997).
[6] K. Maruyama, F. Nori, and V. Vedral, Rev. Mod. Phys. 81, 1 (2009).
[7] T. Sagawa and M. Ueda, Phys. Rev. Lett. **100**, 080403 (2008).
[8] T. Sagawa and M. Ueda, Phys. Rev. Lett. **102**, 250602 (2009).
[9] J. J. Park, K.-H. Kim, T. Sagawa and S. W. Kim, Phys. Rev. Lett. **111**, 230402 (2013).
[10] L. Szilard, Z. Phys. **53**, 840 (1929).
[11] K. Funo, Y. Watanabe and M. Ueda, Phys. Rev. A **88**, 052319 (2013).
[12] S. Popescu, A. J. Short, and A. Winter, Nat. Phys. **2**, 754 (2006).
[13] F. G. S. L. Brandao and M. B. Plenio, Nat. Phys. **4**, 873 (2008).
[14] S. Lloyd, V. Chiloyan, Y. J. Hu, S. Huberman, Z. W. Liu, and G. Chen, arXiv:1510.05035.
[15] S. Toyabe, T. Sagawa, M. Ueda, E. Muneyuki, and M. Sano, Nat. Phys. **6**, 988 (2010).
[16] J. M. R. Parrondo, J. M. Horowitz and T. Sagawa, Nat. Phys. **11**, 131-139 (2015).
[17] F. G. S. L. Brandao, M. Horodecki, H. H. Y. Ng, J. Oppenheim, and S. Wehner, Proc. Natl. Acad. Sci. U.S.A. **112**, 3725 (2015).
[18] J. Oppenheim and S. Wehner, Science **330**, 1072 (2010).
[19] E. Hanggi and S. Wehner, Nat. Commun. 4, 1670 (2013).
[20] L. H. Ren and H. Fan, Phys. Rev. A **90**, 052110 (2014).
[21] M. A. Nielsen, and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000).
[22] M. Berta, M. Christandl, R. Colbeck, J. M. Renes and R. Renner, Nature **6**, 659-662 (2010).
[23] S. Liu, L. Z. Mu, and H. Fan, Phys. Rev. A 91, 042133 (2015).
[24] W. Heisenberg, Z. Phys. **43**, 172 (1927).
[25] Some detailed calculations are presented in the supplementary material.
[26] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, Rev. Mod. Phys. **84**, 1655 (2012).
[27] M. Horodecki, J. Oppenheim, and A. Winter, Nature **436**, 673 (2005).
[28] M. L. Hu and H. Fan, Phys. Rev. A **87**, 022314 (2013).
[29] A. Streltsov, E. Chitambar, S. Rana, M. N. Bera, A. Winter, and M. Lewenstein, Phys. Rev. Lett. **116**, 240405 (2016).

I. SUPPLEMENTARY MATERIAL: SECOND LAW OF THERMODYNAMICS WITH QUANTUM MEMORY

Detailed calculation of the entropy change between initial state and the final state in the heat engine.— We can extract the final state as

$$\rho_{SR}^{(f)} = \mathrm{Tr}_{AB}[\rho^{(f)}] = \sum_k p_k U_{SR}^k \rho_{SR}^{(2)k} U_{SR}^{k\dagger} \tag{25}$$

Considering the concavity of the von Neumann entropy, we have

$$S[\rho_{SR}^{(f)}] \geq \sum_k p_k S[\rho_{SR}^{(2)k}] \tag{26}$$

By means of taking partial trace like equation (25), we can easily obtain $\rho_A^{(f)}$, $\rho_B^{(f)}$ and $\rho_{AB}^{(2)}$, and the results are listed as,

$$S[\rho_A^{(f)}] = S[\rho_A^{(2)}], \qquad S[\rho_B^{(f)}] = S[\rho_B^{(2)}], \tag{27}$$

$$S[\rho_{AB}^{(2)}] = H(p_k) + \sum_k p_k S[\rho_B^{(2)k}]. \tag{28}$$

We then substitute equations (26), (27) and (28) into the relation (10). It is easily to present

$$\begin{aligned} S[\rho_{SR}^{(i)}] - S[\rho_{SR}^{(f)}] &\leq S[\rho_{SR}^{(i)}] - \sum_k p_k S[\rho_{SR}^{(2)k}] \\ &\leq H(p_k) + \sum_k p_k S[\rho_B^{(2)k}] - S[\rho_{AB}^{(i)}] \\ &= S[\rho_{AB}^{(2)}] - S[\rho_{AB}^{(i)}] \\ &\equiv \triangle S_A + \triangle S_B - \triangle I, \end{aligned} \tag{29}$$

where $\triangle S_A = S[\rho_A^{(f)}] - S[\rho_A^{(i)}]$ and I denotes quantum mutual information, and $\triangle I = I(A^{(2)} : B^{(2)}) - I(A^{(i)} : B^{(i)})$ represents the change of quantum mutual information in the process of heat engine. This is the result in the main text.

Work gained from the heat engine by using entropic uncertainty relation.— The entropic uncertainty relation is written as,

$$S(K|B) + S(M|B) \geq \log_2 \frac{1}{c} + S(A|B). \tag{30}$$

Here $S(\cdot|\cdot)$ represents the conditional von Neumann entropy, M, K represent different measurement bases, and c is determined by the overlap between these two bases.

For the heat engine studied by us, if there are two sets of measurement operators, the total work extractable after the heat engine process will be limited due to the uncertainty principle.

The engine will work twice with two sets of measurement operator $\{\Pi_A^k : |k\rangle\langle k|\}$ and $\{\Pi_A^m : |m\rangle\langle m|\}$. Therefore, for state ρ_{AB}^i we have the following relation due to the limitation of the uncertainty principle,

$$S(k|B) + S(m|B) \geq \log_2 \frac{1}{c} + S(A|B), \tag{31}$$

in which $S(A|B) = S(\rho_{AB}^i) - S(\rho_B^i)$. $S(k|B)$ is the conditional von Neumann entropy for the post-measurement state,

$$\begin{aligned} S(k|B) &= S[\sum_k (\Pi_A^k \otimes I_B)\rho_{AB}^i(\Pi_A^k \otimes I_B)] - S(\rho_B^i) \\ &= S[\sum_k p_k|k\rangle\langle k| \otimes \rho_B^{(i)k}] - S(\rho_B^i), \end{aligned} \tag{32}$$

where $p_k = tr(\Pi_A^k \rho_{AB}^i \Pi_A^k)$ and $\rho_B^{(i)k} = tr_A(\Pi_A^k \rho_{AB}^i \Pi_A^k)/p_k$. Similarly,

$$S(m|B) = S[\sum_m q_m|m\rangle\langle m| \otimes \rho_B^{(i)m}] - S(\rho_B^i), \tag{33}$$

where $q_m = tr(\Pi_A^m \rho_{AB}^i \Pi_A^m)$ and $\rho_B^{(i)m} = tr_A(\Pi_A^m \rho_{AB}^i \Pi_A^m)/q_m$. We then substitute these equalities into relation (19). It is easily to obtain

$$\begin{aligned} & S(\sum_k p_k|k\rangle\langle k| \otimes \rho_B^{(i)k}) - S(\rho_B^i) \\ & + S(\sum_m q_m|m\rangle\langle m| \otimes \rho_B^{(i)m}) - S(\rho_B^i) \\ \geq\ & \log_2 \frac{1}{c} + S(\rho_{AB}^i) - S(\rho_B^i) \end{aligned} \tag{34}$$

With working process of the heat engine, we find, $\rho_{AB}^{(2)} = Tr_{SR}\rho^{(2)} = \sum_k p_k|k\rangle_A\langle k| \otimes \rho_B^{(2)k}$. And according to the conclusion of the partial trace, $\rho_B^{(i)k} = tr_A(\Pi_A^k \rho_{AB}^i \Pi_A^k)$ and $\rho_B^{(2)k} = tr_{SR}[tr_A(\Pi_A^k \rho^{(1)\prime} \Pi_A^k)] = tr_A(\Pi_A^k \rho_{AB}^i \Pi_A^k)$. Then we have $\rho_B^{(i)k} = \rho_B^{(2)k}$. Thus the relation (34) can be simplified as

$$S(\rho_{AB}^{(2)}) + S(\rho_{AB}^{(2)\prime}) \geq \log_2 \frac{1}{c} + S(\rho_{AB}^i) + S(\rho_B^i). \tag{35}$$

This equation is the one in the main text.